

# COMPANY OVERVIEW

RELENTLESS  
MEDICINE  
DISCOVERY.





# TANGRAM

## THERAPEUTICS

[+] INTRODUCTION TO TANGRAM TX


# Combining computation and RNAi to solve traditional R&D issues



[+] Industry Problem


[+] Tangram Solution

**Same Targets**



AI-driven novel target ID platform

**High Risk**



Hepatocyte-specific, safe GalOmic siRNAs

**Too Slow**



In silico predictions and accelerated siRNA candidate generation

[+] INTRODUCTION TO TANGRAM TX

# GalOmic RNAi platform enables selective and effective gene silencing.

[+] SAFE

Specific gene silencing in single cell type in the liver limits unwanted side effects

[+] LOW TREATMENT BURDEN

At least quarterly subcutaneous dosing regimen expected in humans ( $\leq 4$  doses/year)

[+] LOW RISK


Commercial-stage modality with large body of safety data and proven high probability of success in clinic

[+] IP PROTECTED


Proprietary GalNAc linkers and GalOmic modification patterns

[+] RNAi mechanism of action


siRNA




Target mRNA



Specific binding




Target gene silenced




[+] INTRODUCTION TO TANGRAM TX

# Liver targeting unlocks a range of opportunities



— The liver interacts with a wide variety of tissues, allowing us to treat a broad range of diseases with our hepatocyte-specific GalOmic medicines



[+] GalOmic Disease  
Area Opportunities


- ▶ Metabolic
- ▶ Hematology
- ▶ Rare disease
- ▶ Cardiovascular
- ▶ Immunology
- ▶ Gastrointestinal
- ▶ Hepatology

[+] MORE

[+] INTRODUCTION TO TANGRAM TX


# We apply AI to uncover differentiated targets.

Realizing hidden opportunities that can make a difference to patients by identifying novel targets backed by robust genetic and biological evidence



[+] INTRODUCTION TO TANGRAM TX

# Our rapid and reproducible process.





[+] INTRODUCTION TO TANGRAM TX

# Our pipeline.

Therapeutic Area	Indication	Discovery	Preclinical	Non-clinical	Clinical
Cardiometabolic	TGM-312	Phase 1/2 ongoing			
	TGM-291				
	TGM-258*				
Hematology	TGM-148	CTA 2026			
Immune-mediated	TGM-407*				
	TGM-462				

\*Partnering opportunity

# TGM-312 FOR MASH



[+] TGM-312 FOR MASH

TGM-312 is a potential safe and effective treatment for MASH, a highly prevalent disease with significant unmet need and opportunity.

>250M

people living with MASH globally, with diagnosis rates increasing



Recent approvals have improved treatment options for patients and **increased diagnosis rates.**



Many patients treated with approved or emerging therapies do not achieve clinically meaningful outcomes.




TGM-312 has potential as a safe and effective MASH treatment with a quarterly subcutaneous dosing regimen.




AM TACUGGGA  
 GUACACUGGGA  
 AM TANGRAM  
 AM TACUGGGA  
 AM TANGRAM  
 ACACUGGG  
 TANGRAM  
 ACACUGGG  
 ACACUGGG  
 ACACUGGG  
 TANGGGG  
 ACACUGGG  
 TANGRAM U  
 TACUGGGGAU  
 ACACUGGGGAU  
 TANGRAM T  
 TACUGGGGAU  
 TANGRAM T  
 ACACUGGGGAU  
 TANGRAM T  
 ACACUGGGGAU  
 ACACUGGGGAU  
 ACACUGGGGAU  
 TANGGGGAU  
 ACACUGGGGAU

[+] TGM-312 FOR MASH


TGM-312 target knockdown supports quarterly dosing regimen in patients



[+] KNOCKDOWN IN MOUSE



[+] KNOCKDOWN IN NHP




[+] TGM-312 FOR MASH


# TGM-312 dramatically improves NAFLD activity scores (NAS) in GAN-DIO MASH mice.

Unprecedented improvement observed with TGM-312 alone and in combination with approved and emerging treatments

[+] STUDY 1 NAS



[+] STUDY 2 NAS



NAFLD activity scores characterise steatosis, inflammation, and ballooning in the liver.

Study 1: 34-week Gubra Amylin NASH diet induction. NAS histology scores measured at 12 weeks post-first dose.


Study 2: 32-week Gubra Amylin NASH diet induction. NAS histology scores measured at 16 weeks post-first dose.

GLP-1R: GLP-1 receptor agonist | THR-beta: thyroid hormone receptor beta agonist | FGF21: FGF21 analogue | GLP-1R/GIPR: dual GLP-1 receptor and GIP receptor agonist


[+] TGM-312 FOR MASH

# Liver histology showcases significant impact of TGM-312 on steatosis and inflammation.

[+] STUDY 1



[+] STUDY 2




Study 1: 34-week Gubra Amylin NASH diet induction. Histology at 12 weeks post-first dose. | Study 2: 32-week Gubra Amylin NASH diet induction. Histology at 16 weeks post-first dose.

GLP-1R: GLP-1 receptor agonist | THR- $\beta$ : thyroid hormone receptor beta agonist | FGF21: FGF21 analogue | GLP-1R/GIPR: dual GLP-1 receptor and GIP receptor agonist


[+] TGM-312 FOR MASH

TGM-312 significantly decreases markers of fibrosis in GAN DIO-MASH mice




— TGM-312 demonstrates equivalent efficacy to emerging anti-fibrotic treatments, FGF21 analogues and GLP-1/GIP receptor agonists.

[+] COLLAGEN 1A1



[+] ALPHA SMOOTH MUSCLE ACTIN




GLP-1R: GLP-1 receptor agonist | THR-beta: thyroid hormone receptor beta agonist | FGF21: FGF21 analogue | GLP-1R/GIPR: dual GLP-1 receptor and GIP receptor agonist

[+] TGM-312 FOR MASH

# Phase 1/2 RESTORE-MASH trial ongoing.

Initial data anticipated 2H 2026



## RESTORE — MASH

[+] PRIMARY ENDPOINTS

Safety, tolerability, and pharmacokinetics/pharmacodynamics

[+] EXPLORATORY ENDPOINTS

Histology, non-invasive biomarkers (MRI-PDFF, Fibroscan), circulating biomarkers

[+] TGM-312 FOR MASH

# Summary and upcoming milestones.




Phase 1/2 RESTORE-MASH trial ongoing with initial safety data anticipated 2H 2026



Results from GAN DIO-MASH studies demonstrate significant potential of TGM-312 as a MASH treatment across patient segments (F2/3 to F4c)




Outstanding preclinical results as monotherapy and in combination with key MOAs across steatosis, inflammation and fibrosis



TGM-312 is a first-on-target, patient-friendly MASH candidate with the potential to drive superior therapeutic outcomes

# TGM-148 FOR BLEEDING DISORDERS



[+] TGM-148 FOR BLEEDING DISORDERS

TGM-148 is a potential pan-bleeding disorder treatment with low treatment burden and low thrombotic risk.

8M

people with bleeding disorders globally, although prevalence of individual disorders are rare.



People with bleeding disorders need prophylactic treatments to prevent bleeds and improve quality of life.



Many treatment regimens remain burdensome and can increase thrombotic risk.




TGM-148 has potential as a pan-bleeding disorder treatment with a desirable safety profile and quarterly subcutaneous dosing.





[+] TGM-148 FOR BLEEDING DISORDERS

TGM-148 normalizes bleeding phenotype in Von Willebrand Factor knockout mice.



[+] ROTATIONAL ELASTOMETRY



Clot formation time normalized by TGM-148



\*\*p<0.01

[+] THROMBIN GENERATION

Thrombin generation parameters normalized by TGM-148




\*p<0.05 | \*\*p<0.01 | \*\*\*\* p<0.0001 | #p<0.05

[+] TGM-148 FOR BLEEDING DISORDERS


# TGM-148 protects knee joints after injury in Hemophilia A and B mice.

[+] NON-INJURED KNEE




- ▶ OARSI Grade 0
- ▶ Defined joint morphology with clear cartilage plate and femoral head

[+] INJURED KNEE + VEHICLE



- ▶ OARSI Grade 5
- ▶ No joint morphology
- ▶ High levels of inflammation
- ▶ Diffuse hemorrhage


[+] INJURED KNEE + TGM-148



- ▶ OARSI Grade 2
- ▶ Defined joint morphology
- ▶ Minor resolving hemorrhage and limited inflammation


[+] TGM-148 FOR BLEEDING DISORDERS

# TGM-148 is safe when administered to wildtype mice with functional clotting



— TGM-148 was safely administered to wildtype mice at exaggerated doses to study end (Day 98). No wildtype mice given exaggerated doses of Qfitlia survived beyond 14 days due to evidence of thrombotic events.

TGM-148 was safely administered to wildtype mice at suprathreshold doses until study end, supporting a desirable safety profile.




Qfitlia dosing led to thrombotic events in wildtype mice, reflecting the narrow safety window observed in clinical trials.

- TGM-148 10 mg/kg
- TGM-148 30 mg/kg
- ▲ Qfitlia 10 mg/kg
- ▼ Qfitlia 30 mg/kg


TGM-148 and Qfitlia administered QW at suprathreshold doses in wildtype mice. Qfitlia doses selected based on Sehgal et al (2015).

[+] TGM-148 FOR BLEEDING DISORDERS


TGM-148 target knockdown supports quarterly dosing regimen in patients



[+] NHP PLASMA KNOCKDOWN



[+] MODELLING OF HUMAN PROFILE



[+] TGM-148 FOR BLEEDING DISORDERS


# Summary and upcoming milestones.




TGM-148 normalizes bleeding phenotype in multiple genetic models of bleeding disorders with duration of action supporting a quarterly dosing regimen in patients



Clean safety profile with evidence of low thrombotic risk in all preclinical and IND-enabling studies to date





CTA submission planned during 2026 with Von Willebrand Disease as the entry indication.





TGM-148 is a first-on-target pan-bleeding disorders candidate with the potential to safely reduce bleeding rates with only four doses a year

# EARLIER PROGRAMS



[+] EARLIER PROGRAMS (OPHTHALMOLOGY)

# TGM-407 is a potential effective, subcutaneous treatment option for Dry AMD.






TGM-407 dosed Day 0 and Day 28.  
Eye protein levels measured Day 56.

## [+] Summary

Preclinical data shows target knockdown in the liver with TGM-407 is tightly correlated with reduced protein expression in the eye, where genetic evidence indicates the target drives Dry AMD pathology. This supports TGM-407 as a potential patient-friendly subcutaneous treatment option with superior efficacy vs current intravitreally administered treatments.

[+] EARLIER PROGRAMS (CARDIOVASCULAR)

# TGM-258 aims to provide a long-acting, safe and effective treatment option for heart failure.







Mice underwent LAD ligation or sham at Day 0. TGM-258 administered QW from Day 0. MRA administered via diet (1g/kg). Functional measurements taken Day 42. MRA: mineralocorticoid receptor antagonist | MI: myocardial infarction

## [+] Summary

TGM-258 improves cardiac function, structure, and serum markers in three distinct models of heart failure, supporting its potential as an effective and patient-friendly treatment option for people living with heart failure. TGM-258 did not lead to adverse events seen with comparator arms in models reflective of clinical experience leading to unmet need.

[+] EARLIER PROGRAMS (CARDIOVASCULAR)

# TGM-291 is a potential broad-acting treatment for cardiometabolic disease.





APOE\*3L.CETP mice on high fat and fructose diet dosed with TGM-291.  
QUICKI: Quantitative Insulin Sensitivity Check Index

## [+] Summary


TGM-291 improves multiple cardiometabolic disease drivers in a humanized metabolic syndrome mouse model and is supported by human genetic evidence linking a missense mutation in the target gene to reduced cardiovascular risk. TGM-291's broad impact across disease drivers beyond LDL-C lowering highlights its potential as a more effective treatment for cardiometabolic diseases.

# LLIBRA OS




[+]LLIBRA OS

# LLibra OS identifies targets and drug candidates for preclinical development.



[+] LLIBRA OS

# LLibra OS enables AI-enhanced target discovery and evaluation.




[+] LLIBRA OS

# Using network analytics and generative AI to identify differentiated targets.

## [+] LLibra Discover


Graph theory approaches including proprietary network analytics and knowledge graph



Analysis of cell-specific or cell-agnostic biological networks and knowledge graphs to identify or validate gene targets.

## [+] LLibra Explore

Generative AI platform with natural language interface and agentic retrieval-augmented generation




Hypothesis generation and validation through exploration of diverse datasets and literature.

[+] LLIBRA OS


# LLibra Evaluate supports portfolio decisions.

Evaluation module assesses therapeutic potential, developability, and commercial viability of hypotheses



[+] LLIBRA OS

# Proprietary siRNA design model accelerates GalOmic medicine design



## [+] PROPRIETARY

Machine learning model trained on thousands of in-house GalOmic datasets

## [+] PREDICTIVE


>80% of leads identified are in the Top 54 sequences

## [+] GENERALIZABLE

Model validated on external patent and clinical leads using unseen constructs

## [+] ACCELERATED

Model enables us to bypass in vitro screening, accelerating discovery phase




Model automatically designs and ranks siRNA sequences

[+] LLIBRA OS

# Our enabling data foundation.

LLibra OS houses a robustly curated data foundation within a flexible, modular architecture



## [+] COMPANY HIGHLIGHTS

# Tangram Therapeutics is poised to unlock the true promise of RNAi medicines.

**Near-term clinical milestones:**

Initial safety data from Phase 1/2 RESTORE-MASH trial of TGM-312 anticipated 2H 2026, CTA submission for TGM-148 anticipated during 2026.

**Validated modality, novel biology:**

Pipeline of differentiated GalOmic medicines with strong preclinical data packages.

**Pragmatic AI deployment:**

State-of-the-art LLibra OS enables continued evolution of our pipeline, science and therapeutic strategy.

**Accelerated, capital-efficient execution:**

Candidate generation in under 12 months, driven by enabling platforms and proven leadership.

# TANGRAM

THERAPEUTICS

[+] HQ

4 Kingdom Street  
Paddington Central  
London W2 6BD

[+] CONTACT

[www.tangramtx.com](http://www.tangramtx.com)  
[contact@tangramtx.com](mailto:contact@tangramtx.com)  
Tel: +44 (0)20 4551 8888

